GRANT COUNTY: Key Data Points

- 4 of the 5 infant deaths in Grant county were neonatal deaths
- A significantly lower percentage of women in Grant county are utilizing prenatal care during the first trimester, 70.8% versus 80.6% respectively.
- A significantly higher percentage of women in the county are smoking during pregnancy, compared to the overall state percentage, 24.3% versus 18.5% respectively.
- The percentage of women breastfeeding on discharge in Grant county is less than the state percentage, 57.3% versus 63.2% respectively.
- Grant county observed 30 accidental deaths (16 of which were motor vehicle), 10 suicidal deaths and 6 homicide

Note: All figures relate to 2003 data from the Indiana State Department of Health, unless otherwise specified

Figure 1. Births

| | То | tal | Wh | ite | Bla | ıck | Oth | ner |
|-----------------|--------|----------------|--------|----------------|--------|----------------|--------|----------------|
| | Number | Crude Rate* | Number | Crude Rate* | Number | Crude Rate* | Number | Crude Rate* |
| Grant County | 799 | 11.2 | 725 | 11.2 | 67 | 13.2 | 7 | 4.2(U) |
| INDIANA | 86,382 | 13.9 | 75,422 | 13.7 | 9,288 | 17.5 | 1,672 | 10.6 |

^{*} births per 1,000 population; U = unstable rate, based on fewer than 20 births

Figure 2. Mortality

| | | NUMBER | | MBER | | RATE* | | |
|-----------|-------------------|--------|------------------------------|-----------------------------|----------------------------|------------------------------|-----------------------------|----------------------------|
| COUNTY OF | REPORTED RACE/ | Live | Infant Deaths (< 1 yr) | Neonatal Deaths (< 28 days) | Post Neonatal Deaths | Infant Deaths (< 1 yr) | Neonatal Deaths (< 28 days) | Post Neonatal Deaths |
| RESIDENCE | ETHNICITY | Births | , ,, | , ,, | (28-364 days) | , ,, | , ,, | (28-364 days) |
| GRANT | All Races | 799 | 5 | 4 | 1 | ** | ** | ** |
| | White | 725 | 4 | 3 | 1 | ** | ** | ** |
| | Black | 67 | 1 | 1 | 0 | ** | ** | 0 |
| | Other | 7 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Hispanic | 35 | 0 | 0 | 0 | 0 | 0 | 0 |
| INDIANA | All Races | 86,382 | 641 | 417 | 224 | 7.4 | 4.8 | 2.6 |
| | White | 75,422 | 479 | 309 | 170 | 6.4 | 4.1 | 2.3 |
| | Black | 9,288 | 148 | 98 | 50 | 15.9 | 10.6 | 5.4 |
| | Other | 1,672 | 14 | 10 | 4 | ** | ** | ** |
| | Hispanic | 6,763 | 38 | 22 | 16 | 5.6 | 3.3 | ** |

^{*}rate per 1,000 live births **Numerator is less than 20 and the rate is unstable

Figure 3. Outcome Indicators of Mother, by Race


| Birth Weight During F | | During Pi | regnancy | | | | | |
|-----------------------|--------|----------------|----------------|-----------|------------------------|---------|---------|----------------|
| GRANT | Total | % Low | % Very Low | % Preterm | % PNC 1st | % | % | % Unmarried |
| COUNTY | Births | (< 2500 grams) | (< 1500 grams) | † | Trimester [‡] | Alcohol | Smoked | Parents |
| All Races | 799 | 9.1 | 1.1 | 12.1 | 70.8(S) | 0.3 | 24.3(S) | 50.1(S) |
| White | 725 | 9 | 0.8 | 12.3 | 72.7(S) | 0 | 25.1(S) | 47.2(S) |
| Black | 67 | 11.9 | 4.5 | 11.9 | 50.7 | 3 | 17.9 | 83.6 |

| Birth Weight | | | | During Pregnancy | | | | |
|--------------|--------|----------------|----------------|-------------------------|-------------|---------|--------|----------------|
| | Total | % Low | % Very Low | % Preterm | % PNC 1st | % | % | % Unmarried |
| INDIANA | Births | (< 2500 grams) | (< 1500 grams) | t | Trimester ‡ | Alcohol | Smoked | Parents |
| All Races | 86,382 | 7.9 | 1.4 | 9.9 | 80.6 | 0.6 | 18.5 | 37.1 |
| White | 75,422 | 7.2 | 1.2 | 9.5 | 82.1 | 0.6 | 19.1 | 32.6 |
| Black | 9,288 | 13.3 | 3.3 | 13.3 | 68.2 | 1.1 | 15.2 | 76.3 |

[†] Preterm: less than 37 weeks gestation

Figure 4. Breastfeeding

| GRANT COUNTY | All Races | White | Black |
|------------------------------------|-----------|-------|-------|
| Total Births | 799 | 725 | 67 |
| Breastfeeding on Discharge | 458 | 430 | 21 |
| % Breastfeeding on Discharge | 57.3 | 59.3 | 31.3 |



[†] PNC: Prenatal Care

'S' Significantly different than the state percent

"**' Outcomes for this county are suppressed due to 20 or fewer total births

| INDIANA | All Races | White | Black |
|------------------------------|--------------|--------|-------|
| Total Births | 86,382 | 75,422 | 9,288 |
| Breastfeeding on Discharge | 54,636 | 49,385 | 3,959 |
| % Breastfeeding on Discharge | 63.2 | 65.5 | 42.6 |




Figure 5. HIV/AIDS

| | Total # of AIDS cases reported since 1982 | Total # of HIV cases reported since 1985 | |
|---------------------|---|--|---|
| GRANT COUNTY | 63 | 2 | 4 |
| INDIANA | 7,641 | 3,85 | 6 |

^{*}numbers current through June 2005

Figure 6. Nutritional Risk Factors by Status,

Indiana WIC Clinics, Quarter Ending 09/30/2004

| Local Agency: Indiana Health Centers, Inc. | | | | | | | | |
|--|--|----------|------------|----------|-----------|-----------|------------------|--|
| Counties Inc. | luded: I | Howard, | Cass, Gran | t, Hamil | ton, Tipt | ton, Clin | iton | |
| | Childr | en 1-5 y | ears of Ag | e | | | % | |
| Risk | | | | | | Total | within | |
| Factor or | White | Black | Hispanic | Asian | Other | | Agency | |
| Status | | | | | | | | |
| Overweight | 112 | 17 | 71 | 0 | 4 | 204 | 4.6% | |
| At Risk for becoming Overweight | 151 | 23 | 64 | 0 | 6 | 244 | 5.5% | |
| Overweight | | | | | | | | |
| | Local Agency: State Wide Counties Included: All | | | | | | | |
| Children 1-5 years of Age | | | | | | | % | |
| Risk Factor or Status | White | Black | Hispanic | Asian | Other | Total | within Agency | |
| Overweight | 1,732 | 445 | 800 | 12 | 97 | 3,086 | 4.8% | |
| At Risk for | | | | | | | | |

| Priority Issues (Grant & Indian | na) | |
|--|---------|---------|
| Demographics: Percentage of population enrolled in Medicaid, Percentage of population enrolled in Food Stamps | Grant | Indiana |
| 2 % of Population enrolled in Medicaid as of June 2003 | 16.2 | 12.4 |
| 3 % of Population in Food Stamps Program, 2004 | 12.0 | 8.6 |
| Priority #1 (pregnancy outcomes): Average Infant Mortality Rate, Average Fetal death Rate | Grant | Indiana |
| 1 Av Total Infant Death Rate (1993-2002) | 9.7 | 8.1 |
| 2 Average Fetal Death Rate (1994-2003) | 7.6 | 6.7 |
| Priority #2 (reduce barriers): Health Professional Shortage Areas, Percentage of women with prenatal care in first trimester | Grant | Indiana |
| Scores for Primary Care Health Professional Shortage Areas (HPSAs) (2005) | 5 | NA |
| 2 % PNC 1st Trimester, 2003 | 70.8 | 80.6 |
| Priority #4 (environmental): Prevalence of asthma, Childhood lead poisoning | Grant | Indiana |
| Prevalence of Asthma in a Medicaid Cohort of Children age 0-17 years, 2003 | 11.2 | 10.4 |
| 2 %Estimated Lead Blood Levels (%EBLL) for Children under 6 years old FY 2004-2005 | 2.4 | 2.1 |
| Priority #5 (tobacco): Percentage of mothers who smoked during pregnancy, Percentage of Adult Smoking | Grant | Indiana |
| 1 % of mothers Smoked During Pregnancy, 2003 | 24.3 | 18.5 |
| 2 % of Adult Smoking in 2003 | 31.9 | 29.3 |
| Priority #7 (adolescent risk behaviors): Adolescent deaths due to motor vehicle accidents, Chlamydia rates (ages 15-24) | Grant | Indiana |
| Number of Adolescent (15-24) deaths due to Motor vehicle accidents (2001-2003) | 9 | 748 |
| 2 Chlamydia rates per 100,000 county specific population (15-24yrs of Age) | 1069.9 | 1541.49 |
| Priority #8 (obesity): Obesity in pregnant women | Grant | Indiana |
| Number of Obese Pregnant Women In Indiana, 2003 | 6 | 2,496 |
| Priority #9 (domestic violence, childhood injury): Child deaths due to Injury, Child abuse and neglect rates | Grant | Indiana |
| 1 Child deaths due to Injury 1-14yrs (2001-2003) | 2 | 257 |
| 2 Child Abuse and Neglect per 1,000 Population under age 18, 2004 | 19.4 | 13.5 |
| Priority #10 (To improve racial and ethnic disparities in women of childbearing age, mothers, and children's health outcomes): Percentage of Hispanic Population of any Race, LBW Disparity Ratio | Grant | Indiana |
| | | |
| 1 % of Hispanic Population of any Race, July,1,2004 2 Low Birth Weight Disparity Ratio (B/W) (1993-2002) | 2.7 | 4.3 |

Source: Data Analysis Team, ISDH. **Notes:** 1. NS= Not Stable 2. NA = Not Available 3. Infant Mortality rate and Fetal Death rate is for 1000 live births. 4. Low Birth Weight: Birth Weight Less than 2,500grams. 5. HPSA Score ranges 1 – 25. The higher the score, the higher the need. 6. EBLL: Ten (10) micrograms of lead per deciliter of whole blood for one venous test; %EBLL is based on screening rate of 27% Statewide, unduplicated kids 7. Obesity in Pregnant Women determined by "Concurrent Illness Obesity" checkbox on the birth certificate 8. PNC percentages are calculated using total births in each county.

Note: The following priorities have been identified for Grant County:

- **Priority # 1:** To decrease high-risk pregnancies, fetal death, low birth weight, infant mortality racial and ethnic disparities in pregnancy outcomes.
- **Priority # 2:** To reduce barriers to access to health care, mental health care and dental care for pregnant women, infants, children, children with special health care needs, adolescents, women and families.
- **Priority # 4:** To reduce morbidity and mortality rates from environmentally related health conditions including asthma, lead poisoning and birth defects.
- **Priority # 5:** To decrease tobacco use in Indiana.
- **Priority # 10:** To improve racial and ethnic disparities in women of childbearing age, mothers, and children's health outcomes.